

CLASSIFICATION: ~~TOP SECRET~~

FOR OFFICIAL USE ONLY

REPORT

STAT

329025

DATE DISTR. 10 August 1948

NO. OF PAGES 1

NO. OF ENCLS.
(LISTED BELOW)

SUPPLEMENT TO
REPORT NO.

CENTRAL INTELLIGENCE AGENCY
~~RESTRICTED~~
INFORMATION REPORT

UNCLASSIFIED

JAN 31 1955

444 / FBD

FOR OFFICIAL USE ONLY

COUNTRY COPY

SUBJECT Scientific Research

PLACE ACQUIRED	USSR
-------------------	------

DATE OF INFORMATION 1946

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE ACT 50 U.S.C. 21 AND 22. AS ASSIGNED, ITS TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW. REPRODUCTION OF THIS FORM IS PROHIBITED. NO OTHER INFORMATION CONTAINED IN BODY OF THE FORM MAY BE UTILIZED AS DEEMED NECESSARY BY THE RECEIVING AGENCY.

THIS IS UNEVALUATED INFORMATION FOR THE RESEARCH
USE OF TRAINED INTELLIGENCE ANALYSTS

SOURCE

Documentary as indicated. (Information specifically requested.)

RECENTLY PUBLISHED RESEARCH OF THE
ALL-UNION INSTITUTE OF ELECTROTECHNOLOGY, USSR

"Organosilicon Compounds: V. Synthesis of Alkyl- and Arylhalomonosilanes," K. A. Andrianov (All-Union Electrotech Inst, Moscow)

"Zhur Obshch Kham" Vol 16, 1946, pp 487-92

Mg was treated with several drops of $(EtO)_4Si$ and then dropwise with $EtBr$, after which more $EtBr$ was added in benzene solution, and mixture was refluxed to yield $EtMgBr$ solution. This was slowly treated with $SiCl_4$ in benzene and refluxed to yield Et_2SiCl_2 . Similarly $iso-BuCl$ gave $iso-BuSiCl_3$; $iso-AmBr$ gave $iso-AmSiCl_3$; use of hexyl bromide gave $C_6H_{13}SiCl_3$; $PhCH_2Cl$ gave $PhCH_2SiCl_3$ in a similar procedure. $1-C_6H_5MgBr$ and Mg moistened with a little $(EtO)_4Si$ in 2 volumes benzene was heated until the Grignard reagent was prepared and added to $SiCl_4$ and heated on a steam bath; $MgClBr$ was filtered off and the original layer was distilled in vacuo to yield $1-C_6H_5SiCl_3$. Preparation of dialkyl derivatives was conducted as above with amount of $SiCl_4$ being halved, i.e., $2MgBr + SiCl_4$. Thus, the following were prepared: Et_2SiCl_2 ; $(C_6H_5)_2SiCl_2$; $iso-Bu_2SiCl_2$.

- END -

- 1 -

CLASSIFICATION

FOR OFFICIAL USE ONLY

[illegible]

RECEIVED